



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Mr. C. F. GUMMER solved the problem in a similar manner and added:

It may be of interest to find the expectation. This will be

$$\sum_{r=k}^{r=N-n+k} \frac{\binom{r-1}{k-1} \binom{N-r}{n-k}}{\binom{N}{n}} r = \frac{k}{N} f(N, n, k),$$

where r is the number on the k th ball, and

$$f(N, n, k) = \sum_{r=k}^{r=N-n+k} \binom{N-n}{r-k} \frac{r}{k} \frac{N-r}{n-k}.$$

From the relation

$$\binom{N-n}{r-k} = \binom{N-n-1}{r-k-1} + \binom{N-n-1}{r-k},$$

it follows that

$$f(N, n, k) = (n-k+1)f(N, n+1, k) + (k+1)f(N, n+1, k+1). \quad (1)$$

Now $f(N, N, k) = 1$, being independent of k . Therefore, by (1), $f(N, N-1, k) = N+1$, also independent of k , and finally $f(N, n, k) = (N+1)N \cdots (n+2)$. Hence, the expectation for the k th ball is $k(N+1)/(n+1)$.

2699 [May, 1918]. Proposed by the late ROGER E. MOORE, University of Wisconsin.

Show that if $a_k^{(r)}$ denotes the k th term of an arithmetic progression of order r , and c_k denotes the k th binomial coefficient in the expansion of $(a-b)^n$, n being a positive integer,

$$s \equiv \sum_{k=1}^{n+1} c_k a_k^{(r)} = 0, \quad \text{if} \quad n > r.$$

SOLUTION BY ELBERT H. CLARKE, Hiram College.

Let d_0 be the first term in the arithmetic progression and let d_1, \dots, d_r denote the initial difference of each order. Using the usual abbreviated notation for binomial coefficients, we write

$$a_k^{(r)} = \binom{k-1}{0} d_0 + \binom{k-1}{1} d_1 + \binom{k-1}{2} d_2 + \cdots + \binom{k-1}{r} d_r,$$

$$c_k = (-1)^{k-1} \binom{n}{k-1}$$

and

$$\sum_{k=1}^{n+1} c_k a_k^{(r)} = \sum_{i=0}^r d_i \sum_{k=1}^{n+1} (-1)^{k-1} \binom{k-1}{i} \binom{n}{k-1}.$$

Consider the inner sum. Since $\binom{k-1}{i} = 0$, for $k < i+1$,

$$\sum_{k=1}^{n+1} (-1)^{k-1} \binom{k-1}{i} \binom{n}{k-1} = \sum_{k=i+1}^{n+1} (-1)^{k-1} \binom{k-1}{i} \binom{n}{k-1},$$

and the latter expression easily becomes

$$\binom{n}{i} \sum_{k=i+1}^{n+1} (-1)^{k-1} \binom{n-i}{k-(i+1)}.$$

Now put $k-i-1 = t$ and we have

$$\binom{n}{i} \sum_{t=0}^{n-i} (-1)^{t+i} \binom{n-i}{t}.$$

But the expression under summation is simply $(-1)^i (1-1)^{n-i}$. Hence, the coefficient of every d_i is zero. Therefore,

$$\sum_{k=1}^{n+1} c_k a_k^{(r)} = 0, \quad n > r.$$

NOTE.—This proof is good only for $n > r$. Numerical examples can easily be given to show that the formula is not true for $n \equiv r$.

Also solved by HORACE L. OLSON.

NOTES AND NEWS.

EDITED BY E. J. MOULTON, Northwestern University, Evanston, Ill.

Dr. J. N. VAN DER VRIES has resigned his position as professor of mathematics at the University of Kansas to continue his work as secretary of the central district of the Chamber of Commerce of the United States, with headquarters at Chicago.

Mr. G. H. CRESSE, previously instructor in the University of Michigan, has been appointed to an instructorship in mathematics at the U. S. Naval Academy at Annapolis; he was granted the degree of doctor of philosophy at the University of Chicago in December.

Mr. F. S. NOWLAN, of Bowdoin College, has been promoted to an assistant professorship of mathematics.

Dr. MARION B. WHITE, formerly of the Ypsilanti State Normal College, Michigan, is now professor of mathematics at Carleton College.

Dr. TOBIAS DANTZIG and Dr. G. A. PFEIFFER have been appointed instructors in mathematics at Columbia University.

Professor GRIFFITH C. EVANS is now scientific attaché to the American Embassy at Rome.

Dr. W. GROSS, of the University of Vienna, has been promoted to professor of mathematics.

Professor E. WIECHERT of Göttingen has been appointed professor of geodesy and geophysics at the University of Berlin.

Efforts are being made to establish a chair of mathematical physics at the University of Edinburgh in memory of the late Professor TAIT.

At the University of Strasbourg Professor RENÉ M. FRÉCHET, of the University of Poitiers, has been appointed professor of mathematics, and PIERRE WEISS, professor at the Polytechnikum, Zurich, professor of general physics.

CHARLES L. DOOLITTLE, professor emeritus of astronomy at the University of Pennsylvania, and director of the Flower Observatory, died on March 3, 1919, aged seventy-five years.